

Surname	Centre Number	Candidate Number
First name(s)		2



GCE AS

B410U20-1



S25-B410U20-1



TUESDAY, 20 MAY 2025 – MORNING

CHEMISTRY – AS component 2

Energy, Rate and Chemistry of Carbon Compounds

1 hour 30 minutes

Section A

Section B

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1. to 6.	10	
7.	10	
8.	18	
9.	12	
10.	12	
11.	18	
Total	80	

B410U201
01

ADDITIONAL MATERIALS

- A calculator, pencil and ruler
- **Data Booklet** supplied by WJEC

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use gel pen or correction fluid.

You may use a pencil for graphs and diagrams only.

Write your name, centre number and candidate number in the spaces at the top of this page.

Section A Answer **all** questions.

Section B Answer **all** questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

The maximum mark for this paper is 80.

Your answers must be relevant and must make full use of the information given to be awarded full marks for a question.


The assessment of the quality of extended response (QER) will take place in **Q10(a)**.



JUN25B410U20101

SECTION AAnswer **all** questions.

1. The skeletal formula of a compound is shown below.



Name the compound.

[1]


.....

2. Draw the structure of 2,3-dimethylpentanoic acid.

[1]

3. Name the alkene monomer that can be polymerised to give the following polymer.

[1]



.....




4. (a) State why ammonia is classified as a nucleophilic reagent. [1]

.....

- (b) Write the equation for the standard molar enthalpy change of formation of ammonia. [2]

.....

5. The diagram below shows the distribution of molecular energies for a sample of gas. Use this curve to explain how a catalyst increases the rate of reaction. [2]



.....

.....

.....

6. Calculate the number of **molecules** in 25.6 kg of ethanoic acid (M_r 60.0). [2]

Molecules of ethanoic acid =



SECTION B

Answer **all** questions.

7. (a) A student was given a sample of a halogenoalkane and told to find out which halogen was present in the sample.

His method, observation and conclusion are shown below.

- Pipette 1 cm^3 of the halogenoalkane into a test tube
- Add 1 cm^3 of 1 mol dm^{-3} aqueous sodium hydroxide and warm
- Add 3-4 drops of 1 mol dm^{-3} dilute nitric acid
- Add 1 cm^3 of aqueous silver nitrate
- Cream precipitate is seen so halogen is bromine

- (i) Suggest an improvement to the method that does not require the use of an additional reagent. Give a reason for your suggestion. [2]

.....

.....

- (ii) Another student thought that the precipitate was white and that the halogen present in the sample was therefore chlorine.

Give a further test that the student could carry out to prove which halogen is present in the sample. Include the results for chlorine and bromine in your answer. [2]

.....

.....

.....



- (b) When 6.26 g of 1-bromobutane (M_r 137) was converted into pentanenitrile, C_4H_9CN , 2.58 g of the product was obtained.

Calculate the percentage yield for this reaction. Give your answer to an **appropriate** number of significant figures. [3]

Percentage yield = %

- (c) Despite a total ban on CFCs coming into effect in 2014, many companies are still in the process of replacing their cooling systems. Consequently, there are plenty of air-conditioning systems still using CFCs such as Freon, CHF_2Cl .

Explain how Freon poses a threat to the environment. [3]

.....

.....

.....

.....

.....

.....





8. The information given below relates to a corrosive white solid, **X**.

Quantitative analysis shows that its percentage composition by mass is carbon 33.2%, chlorine 32.7% and hydrogen 4.6% with the remainder being oxygen.

The mass spectrum shows two molecular ion signals at m/z 108 and m/z 110 in the ratio of 3:1.

On addition to an aqueous solution of sodium hydrogencarbonate, effervescence is seen.

Its infrared spectrum and low resolution ^1H NMR spectrum are shown below.



(a) Use **all** the information to identify compound **X**.

[9]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

B410U201
07



- (b) Ethanol is normally produced worldwide either by the hydration of ethene or the fermentation of crops such as sugarcane.

- (i) The hydration of ethene is typically carried out using a catalyst of phosphoric acid at a pressure of 70 atm.

State the temperature used in this reaction.


[1]

..... °C

- (ii) Using the fermentation method, the ethanol is separated by distillation.

Complete and label the diagram of the apparatus used in a laboratory to carry out a simple distillation.

[3]



(c) Ethanol can be used to make many useful compounds.

(i) It can be dehydrated to form ethene.

Name a suitable dehydrating agent for this reaction.

[1]

.....

(ii) It can be oxidised by acidified potassium dichromate(VI) to form ethanoic acid.

State what you would observe during this reaction.

[1]

.....

(iii) I. Classify the type of reaction taking place when ethanol reacts with propanoic acid using sulfuric acid as a catalyst.

[1]

.....

II. Draw the structure of the organic product formed in this reaction.

[1]

(d) Ethanoic acid reacts with sodium hydroxide to form sodium ethanoate and water.

Write an equation for this reaction.

[1]

.....



9. (a) A student was asked to measure the enthalpy change that occurs when sodium nitrate is dissolved in water.

She was told to use the following method.

- Weigh 8.70 g of sodium nitrate
- Accurately measure 50.0 cm³ of water into a polystyrene cup (placed in another polystyrene cup to provide insulation)
- Place a 0.2 °C graduated thermometer in the water and measure its temperature every minute, stirring before reading the temperature
- At the third minute add the sodium nitrate, but do not record the temperature
- Stir the mixture thoroughly, then record the temperature after four minutes
- Continue stirring and record the temperature every minute for a further six minutes

Her results are shown below.

Time /min	0	1	2	3	4	5	6	7	8	9	10
Temp /°C	19.8	20.0	20.0	–	14.8	12.2	12.6	13.0	13.2	13.3	13.4

- (i) Give the reason why she accurately measured the volume of the water. [1]

.....

.....

- (ii) Give the reason why the temperature of the water was measured for two minutes before adding the sodium nitrate. [1]


.....

.....



- (iii) Plot the results on the grid below.

[2]

Examiner
only

- (iv) Determine the maximum temperature **change** by drawing appropriate lines of best fit.

[3]

Maximum temperature change = °C



- (v) Explain why the value determined in part (iv) is a more accurate temperature change than one found using the lowest temperature recorded in the experiment.

[1]

.....

.....

.....

- (vi) The student calculated the enthalpy change to be 19.2 kJ mol^{-1} . The data book value for this enthalpy change is 20.5 kJ mol^{-1} .

Express the difference between these two values as a percentage of the data book value.


[1]

Percentage difference = %



- (b) Sulfur dioxide can be formed in the atmosphere when hydrogen sulfide, emitted in gases from volcanoes, reacts with the air.

The equation and the enthalpy change for the reaction are as follows.



- (i) Use this enthalpy change of reaction and the standard enthalpy changes of formation ($\Delta_f H^\theta$) given in the table below to calculate the standard enthalpy change of formation of sulfur dioxide. [2]

Substance	$\Delta_f H^\theta / \text{kJ mol}^{-1}$
$\text{H}_2\text{S}(\text{g})$	-20.2
$\text{O}_2(\text{g})$	0
$\text{H}_2\text{O}(\text{l})$	-286


$$\Delta_f H^\theta = \dots\dots\dots \text{kJ mol}^{-1}$$

- (ii) Suggest why scientists are concerned about this reaction. [1]

.....



10. (a) Hydrogen peroxide, H_2O_2 , decomposes slowly to give water and oxygen.



If a catalyst such as manganese(IV) oxide is used, the rate of decomposition increases significantly.

A student was asked to determine the initial rate of decomposition of a solution of H_2O_2 at a constant temperature of 40°C using a manganese(IV) oxide catalyst.

Outline how the student should do this.

Your answer should include

- a method for the experiment, including essential apparatus
- a description of how the results would be used to determine the initial rate

[6 QER]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



- (b) State how the overall rate of decomposition would compare with the initial rate of decomposition.

Explain your answer using particle theory.

[2]

.....

.....

.....

.....

- (c) State the type of catalyst used in this experiment.

[1]

.....

- (d) Another student said that lead(IV) oxide is a more efficient catalyst.

Briefly describe how the student could determine which is the better catalyst in the decomposition of hydrogen peroxide.

[3]

.....

.....

.....

.....

.....

.....



11. (a) There are many isomers with the formula C_5H_{10} . Information about five of these isomers, **A**, **B**, **C**, **D** and **E**, is given below.
- **A** and **B** are straight-chain isomers and react with hydrogen and a catalyst to give the same hydrocarbon
 - **B** is the only one that shows *E–Z* isomerism
 - **C** does not react with hydrogen or hydrogen bromide
 - **D** and **E** are branched-chain isomers and react with hydrogen and a catalyst to give the same hydrocarbon
 - **D** and **E** react with hydrogen bromide to form the same bromoalkane as the major product in each case
 - One of the compounds is a cycloalkane



- (i) Use the information to deduce **possible** identities for each isomer. Explain your reasoning. [8]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Name the catalyst in the reaction between isomers **A** and **B** and hydrogen. [1]

.....

- (iii) Classify the reaction mechanism when isomers **D** and **E** react with hydrogen bromide. [1]

.....



- (b) A technician requires a sample of hexane. It is kept in a metal storage cupboard. The bottle on the shelf where it is usually found has an incomplete label.

The technician carries out the following procedure and is satisfied that the liquid is hexane.

She injects a 0.142 g sample of the liquid into a sealed gas syringe and places it in an oven at 383 K. The liquid vaporises. The volume of the gas produced is 52.0 cm³ at a pressure of 101 kPa.

- (i) Show that the technician is correct. [4]

- (ii) Suggest why the bottle of hexane is kept in a metal storage cupboard. [1]

.....



- (iii) Explain how useful ^{13}C NMR spectra and low resolution ^1H NMR spectra would be to distinguish between hexane and pentane. [3]

Examiner
only

.....

.....

.....

.....

.....

.....

18

END OF PAPER



