

Please check the examination details below before entering your candidate information

|                      |                      |                      |                      |                      |                      |                      |                      |                      |  |
|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|--|
| Candidate surname    |                      |                      |                      |                      | Other names          |                      |                      |                      |  |
| Centre Number        |                      |                      |                      |                      | Candidate Number     |                      |                      |                      |  |
| <input type="text"/> | <input type="text"/> | <input type="text"/> | <input type="text"/> | <input type="text"/> | <input type="text"/> | <input type="text"/> | <input type="text"/> | <input type="text"/> |  |

**Pearson Edexcel Level 3 GCE**

**Tuesday 13 May 2025**

Morning (Time: 1 hour 30 minutes)

Paper reference **8CH0/01**

**Chemistry**  
**Advanced Subsidiary**  
**PAPER 1: Core Inorganic and Physical Chemistry**

**You must have:**  
 Scientific calculator, Data Booklet

Total Marks

## Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided  
 – *there may be more space than you need.*

## Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets  
 – *use this as a guide as to how much time to spend on each question.*
- For the question marked with an **asterisk (\*)**, marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other where appropriate.
- A Periodic Table is printed on the back cover of this paper.

## Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P77830A

©2025 Pearson Education Ltd.  
 Y:1/1/1/



  
**Pearson**

Answer ALL questions.

Some questions must be answered with a cross in a box ☐. If you change your mind about an answer, put a line through the box ☒ and then mark your new answer with a cross ☐.

- 1 Chlorine,  $\text{Cl}_2$ , and silicon tetrachloride,  $\text{SiCl}_4$ , are covalently bonded molecules with low boiling temperatures.

(a) The definition of electronegativity is the ability of an atom to attract

(1)

- ☐ **A** a single bonding electron from a covalent bond
- ☐ **B** a pair of electrons from a covalent bond
- ☐ **C** electron density from a neighbouring atom
- ☐ **D** two electrons from a positive ion

- (b) Draw a dot-and-cross diagram of silicon tetrachloride, showing outer shells of electrons only.

Use crosses (x) for silicon electrons and dots (•) for chlorine electrons.

(2)



- (c) Explain, using electronegativity values, why both chlorine and silicon tetrachloride are non-polar molecules.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) Explain why the boiling temperature of chlorine ( $-34^{\circ}\text{C}$ ) is lower than that of silicon tetrachloride ( $57^{\circ}\text{C}$ ).

(2)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**(Total for Question 1 = 8 marks)**

.....



2 At room temperature, both sodium metal and sodium chloride are solids containing ions.

- (a) (i) Draw labelled 2-dimensional diagrams showing the structures of both sodium metal and sodium chloride including **nine** ions in each diagram.

(3)

|              |                 |
|--------------|-----------------|
| sodium metal | sodium chloride |
|--------------|-----------------|

- (ii) Describe what holds the ions together in sodium metal and in sodium chloride.

(2)

Sodium metal

.....

.....

.....

Sodium chloride

.....

.....

.....

- (iii) The melting temperature of sodium chloride is much higher than that of sodium metal.

State what this information shows about the bonding in these two substances.

(1)

.....

.....

.....



- (iv) Compare and contrast the electrical conductivity of sodium metal with that of sodium chloride, including how the charge is carried.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Sodium hydride, which contains the hydride ion,  $\text{H}^-$ , has the same type of structure as sodium chloride.

- (i) Sodium hydride can be made by reacting sodium metal with hydrogen.


Write the equation for this reaction. State symbols are not required.

(1)

- (ii) Sodium hydride reacts with aluminium chloride to produce sodium tetrahydridoaluminate(III),  $\text{NaAlH}_4$ .

Complete a balanced equation for this reaction.

(1)







(iii)  $\text{H} \rightarrow \text{Al}$  represents one of the bonds in  $\text{AlH}_4^-$ .

State the name of this type of bond and how the bond forms in this ion.

(2)

(iv) What are the shapes of the aluminium chloride molecule and the tetrahydridoaluminate(III) ion?

(1)

|                                   | Name of shape of $\text{AlCl}_3$ | Diagram of shape of $\text{AlH}_4^-$                                                 |
|-----------------------------------|----------------------------------|--------------------------------------------------------------------------------------|
| <input type="checkbox"/> <b>A</b> | triangular-based pyramid         |   |
| <input type="checkbox"/> <b>B</b> | triangular-based pyramid         |  |
| <input type="checkbox"/> <b>C</b> | trigonal planar                  |  |
| <input type="checkbox"/> <b>D</b> | trigonal planar                  |  |

(Total for Question 2 = 14 marks)



3 This question is about intermolecular forces.

(a) What are the intermolecular forces between water molecules?

(1)

- ☐ A hydrogen bonds only
- ☐ B London forces and permanent dipoles only
- ☐ C London forces, permanent dipoles and hydrogen bonds
- ☐ D permanent dipoles and hydrogen bonds only

(b) Ethanol,  $\text{CH}_3\text{CH}_2\text{OH}$ , is soluble in water but ethanethiol,  $\text{CH}_3\text{CH}_2\text{SH}$ , is almost insoluble.

The **best** explanation for the difference in solubility is that

(1)

- ☐ A ethanethiol molecules form stronger London forces with water than ethanol
- ☐ B sulfur is less electronegative than oxygen
- ☐ C ethanol forms hydrogen bonds with water but ethanethiol does not
- ☐ D ethanol has a larger dipole than ethanethiol

(Total for Question 3 = 2 marks)

4 This question is about magnesium nitrate,  $\text{Mg}(\text{NO}_3)_2$ .

Magnesium nitrate can be made by the reaction of magnesium metal with nitric acid. A piece of magnesium metal was added to each of two conical flasks containing the same volume of nitric acid.

One flask contains dilute nitric acid and the other flask contains concentrated nitric acid.

In both flasks the acid was in excess.

- (a) In the flask containing dilute nitric acid, the magnesium reacted, producing bubbles of a colourless gas.

(i) Give the name of the gas and the test for the gas, with the positive result.

(2)

- (ii) Write an equation for the reaction that occurs with magnesium and dilute nitric acid.

Include state symbols.

(2)

- (iii) The flask containing concentrated nitric acid also produced bubbles, but this time the gas was nitrogen dioxide. Water was also formed.

Write an equation for the reaction that occurs with concentrated nitric acid. State symbols are not required.


(1)



(b) Magnesium nitrate thermally decomposes, as shown by the equation.



A sample of magnesium nitrate was heated using the apparatus shown. The resultant mixture of gases was passed through water containing universal indicator, removing one of the products. This green solution turned red. The remaining gas was collected by displacement of water.



- (i) State the appearance of the mixture of gases **before** bubbling through the universal indicator solution, and the appearance of the gas **after** being collected.

(1)

- (ii) Explain the change in appearance of the mixture of gases, and of the universal indicator solution.

(2)

- (iii) The measuring cylinder was full of water before a sample of magnesium nitrate was thermally decomposed.  
Part of the measuring cylinder after the reaction had finished is shown.



The volume of gas collected was

(1)

- ☐ **A** 218 cm<sup>3</sup>
- ☐ **B** 236 cm<sup>3</sup>
- ☐ **C** 257 cm<sup>3</sup>
- ☐ **D** 264 cm<sup>3</sup>

- (iv) The sample of magnesium nitrate had a mass of 2.966 g.

Calculate the molar volume of the gas collected, giving your answer to 3 significant figures. Include units in your answer.

You should assume that the decomposition was complete.

The volume of gas was measured at standard temperature and pressure.

Use of  $pV = nRT$  is not required.

[ $M_r$  of magnesium nitrate = 148.3]

(4)



- (v) Explain why **three** significant figures is appropriate for the answer in this calculation.

(2)

(Total for Question 4 = 15 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- 5 The metal zinc (a d-block element) is chemically similar to magnesium (a Group 2 element). One reason for this is because both elements form ionic compounds containing stable  $2+$  ions.

(a) (i) Complete the electronic structure of the Zn atom and of the  $\text{Mg}^{2+}$  ion.

(2)

Zn  $1s^2$  .....

$\text{Mg}^{2+}$   $1s^2$  .....

- (ii) Explain, in terms of ionisation energies, why zinc and magnesium both form stable  $2+$  ions.

(2)

.....

.....

.....

.....

.....

.....

- (b) Naturally occurring samples of zinc have five isotopes, the most abundant being  ${}^{64}_{30}\text{Zn}$ .

(i) Complete the table.

(1)

|                         | Number of protons | Number of neutrons |
|-------------------------|-------------------|--------------------|
| ${}^{64}_{30}\text{Zn}$ |                   |                    |

- (ii) The relative isotopic mass of  ${}^{64}_{30}\text{Zn} = 63.929$ .

Give the reason why the relative isotopic mass is not exactly 64.

(1)

.....

.....

.....



- (iii) The table gives the relative abundances and isotopic masses of four of the five isotopes.

| Symbol of isotope      | $^{64}_{30}\text{Zn}$ | $^{66}_{30}\text{Zn}$ | $^{67}_{30}\text{Zn}$ | $^{70}_{30}\text{Zn}$ |
|------------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Relative isotopic mass | 63.929                | 65.926                | 66.927                | 69.925                |
| Abundance / %          | 48.63                 | 27.90                 | 4.10                  | 0.62                  |

The relative atomic mass of the sample was found to be 65.396.

Calculate the relative isotopic mass of the fifth isotope, giving your answer to 3 decimal places, and hence give the symbol of the fifth isotope.

(4)

Symbol of the fifth isotope .....

- (iv) The mass of an atom of  ${}^{64}_{30}\text{Zn} = 1.06157 \times 10^{-22} \text{ g}$ .

Calculate the value of the Avogadro constant using the mass of this atom and the relative isotopic mass, giving your answer to an appropriate number of significant figures. You **must** show your working.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- (c) Both zinc and magnesium are essential elements for human health, and both can be taken as a dietary supplement in the form of their sulfates. Hydrated zinc sulfate,  $\text{ZnSO}_4 \cdot x\text{H}_2\text{O}$ , can be taken orally as tablets. The value of  $x$  can be found by heating the hydrated zinc sulfate until there is no further change in mass.

450 mg of hydrated zinc sulfate gave 253 mg of anhydrous zinc sulfate on heating.


Calculate the value of  $x$  in  $\text{ZnSO}_4 \cdot x\text{H}_2\text{O}$ .

(4)

(Total for Question 5 = 15 marks)

- 6 When potassium iodide and concentrated sulfuric acid are mixed, several reactions occur.

The equation for one of these reactions is shown.



- (a) (i) Explain how the equation indicates that the reaction should be carried out in a fume cupboard rather than in the laboratory.

(2)

.....

.....

.....

.....

.....

.....

- (ii) Give **two** observations that would be made when the reaction shown by the equation is carried out in a fume cupboard.

(2)

.....

.....

.....

.....

.....

- (b) Which change in oxidation number of sulfur, if any, occurs during the formation of  $\text{H}_2\text{S}$ ?

(1)

- ☐ **A** from +6 to -2
- ☐ **B** from +6 to +2
- ☐ **C** from +2 to -2
- ☐ **D** no change



- (c) Deduce the two ionic half-equations for this reaction.  
State symbols are not required.

(2)

Oxidation half-equation

Reduction half-equation

- (d) When potassium chloride and concentrated sulfuric acid are mixed, misty fumes are observed from the only reaction that occurs.

- (i) Identify, by name or formula, the misty fumes.

(1)

- (ii) Explain what this observation indicates about the relative strength of the chloride ion and the iodide ion as reducing agents.

(2)

(Total for Question 6 = 10 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



7 This question is about periodic properties of elements.

(a) Explain why the atomic radius decreases across Period 3 from sodium to chlorine.

(2)

---

---

---




---

---

---

(b) Which equation does **not** represent the reaction that occurs during the first ionisation energy for a Period 2 element?

(1)




(c) Which isoelectronic ion has the smallest ionic radius?

(1)



\*(d) The graph shows the first ionisation energy values across Period 3.



Explain these changes in first ionisation energies.

(6)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

Handwriting practice area with 20 horizontal dotted lines.

(Total for Question 7 = 10 marks)



- 8 Ferrocene is a compound containing iron, carbon and hydrogen only. The amounts of carbon and hydrogen in a sample can be determined using the apparatus shown.



The hydrogen atoms from the sample react with oxygen to form water, which is absorbed in the first U-tube.

The carbon atoms react with oxygen to form carbon dioxide, which is absorbed in the second U-tube.

A 0.830 g sample of ferrocene was analysed using this apparatus.

The table shows the mass of the U-tubes before and after combustion.

|                                                  |                            |       |
|--------------------------------------------------|----------------------------|-------|
| U-tube containing solid absorbing water          | Mass before combustion / g | 36.34 |
|                                                  | Mass after combustion / g  | 36.74 |
| U-tube containing solid absorbing carbon dioxide | Mass before combustion / g | 35.28 |
|                                                  | Mass after combustion / g  | 37.23 |

Mass spectrometry shows that ferrocene has a molecular ion peak at  $m/z = 186$ .



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Deduce the molecular formula of ferrocene using all of the data.

(6)

---

(Total for Question 8 = 6 marks)

---

**TOTAL FOR PAPER = 80 MARKS**



## 24

P 7 7 8 3 0 A 0 2 4 2 4

\* Lanthanide series

\* Actinide series

|     |    |         |    |     |    |              |    |     |    |           |    |       |    |            |    |     |    |           |    |     |    |           |    |     |    |            |    |     |    |           |    |     |    |             |    |     |    |             |    |     |    |         |     |     |    |           |     |     |    |           |     |     |    |            |     |
|-----|----|---------|----|-----|----|--------------|----|-----|----|-----------|----|-------|----|------------|----|-----|----|-----------|----|-----|----|-----------|----|-----|----|------------|----|-----|----|-----------|----|-----|----|-------------|----|-----|----|-------------|----|-----|----|---------|-----|-----|----|-----------|-----|-----|----|-----------|-----|-----|----|------------|-----|
| 140 | Ce | cerium  | 58 | 141 | Pr | praseodymium | 59 | 144 | Nd | neodymium | 60 | [147] | Pm | promethium | 61 | 150 | Sm | samarium  | 62 | 152 | Eu | europium  | 63 | 157 | Gd | gadolinium | 64 | 159 | Tb | terbium   | 65 | 163 | Dy | dysprosium  | 66 | 165 | Ho | holmium     | 67 | 167 | Er | erbium  | 68  | 169 | Tm | thulium   | 69  | 173 | Yb | ytterbium | 70  | 175 | Lu | lutetium   | 71  |
| 232 | Th | thorium | 90 | 231 | Pa | protactinium | 91 | 238 | U  | uranium   | 92 | 237   | Np | neptunium  | 93 | 242 | Pu | plutonium | 94 | 243 | Am | americium | 95 | 247 | Cm | curium     | 96 | 245 | Bk | berkelium | 97 | 251 | Cf | californium | 98 | 254 | Es | einsteinium | 99 | 253 | Fm | fermium | 100 | 256 | Md | mercurium | 101 | 254 | No | nobelium  | 102 | 257 | Lr | lawrencium | 103 |

**DO NOT WRITE IN THIS AREA**