

ADVANCED General Certificate of Education 2016

 C	Centr	e Nu	mber
Can	didat	e Nu	mber

Chemistry

Assessment Unit A2 1

assessing

Periodic Trends and Further Organic, Physical and Inorganic Chemistry



AC212

[AC212]

FRIDAY 27 MAY, MORNING

TIME

2 hours.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number in the spaces provided at the top of this page.

Answer all sixteen questions.

Answer **all ten** questions in **Section A**. Record your answers by marking the appropriate letter on the answer sheet provided. Use only the spaces numbered 1 to 10. Keep in sequence when answering.

Answer all six questions in Section B.

You must answer the questions in the spaces provided.

Do not write outside the boxed area on each page or on blank pages.

Complete in blue or black ink only. Do not write with a gel pen.

INFORMATION FOR CANDIDATES

The total mark for this paper is 120.

Quality of written communication will be assessed in Question 16(a).

In Section A all questions carry equal marks, i.e. **two** marks for each question.

In Section B the figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

A Periodic Table of Elements, containing some data, is included in this question paper.



Section A

)

ng Learning

)

0

)

)

Rewardin DD

)

0

FE

)

)

)

Rewardin Pag sg Learning

6

For each of the questions only one of the lettered responses (A-D) is correct.

Select the correct response in each case and mark its code letter by connecting the dots as illustrated on the answer sheet.

1 For the reaction,

$$\mathrm{H_2O_2} \, + \, 2\mathrm{I^-} \, + \, 2\mathrm{H^+} \, \longrightarrow \mathrm{I_2} \, + \, 2\mathrm{H_2O}$$

the rate law is given by the equation, rate = $k[H_2O_2][I^-][H^+]$.

Which one of the following is the factor by which the rate of reaction increases if the concentration of each reactant is doubled?

- A 2
- B 4
- C 8
- D 16
- 2 23 mg of sodium were added to 100 cm³ of water. Which one of the following is the pH of the solution produced?
 - A 10
 - B 11
 - C 12
 - D 13



- 3 Which one of the following is the number of isomers with the molecular formula C_4H_9Br ?
 - A 2

Rewarding L.

Learning

GE

Rewarding L

GE

Rewarding L

Rewarding L

Rewarding L

Rowarding L

Leaming

Rewarding L

Rowarding L

Rewarding L

- B 3
- C 4
- D 5
- **4** Which one of the following oxides produces a slightly alkaline solution when dissolved in water?
 - A Al_2O_3
 - B MgO
 - C Na₂O
 - D SiO₂
- **5** Which one of the following does **not** have units?
 - A ΔS for the boiling of water
 - B ΔG for the Haber process
 - C K_w for water at 30 °C
 - $\mathsf{D} \quad \mathsf{K}_{\mathsf{c}} \text{ for the esterification of ethanol} \\$

[Turn over



6 Phosphate ions are hydrolysed by water in the following equilibrium:

$$PO_4^{3-} + H_2O \implies HPO_4^{2-} + HO^{-}$$

)

)

)

Which one of the following shows the Brønsted-Lowry bases in the equilibrium?

- A PO_4^{3-} and HPO_4^{2-}
- B PO_4^{3-} and HO^-
- C H₂O and HO⁻
- D H₂O and HPO₄²⁻
- 7 Reactions which have a large activation energy have a
 - A small entropy change.
 - B large entropy change.
 - C small rate constant.
 - D large rate constant.
- **8** Which one of the following reagents would **not** be used to distinguish between aldehydes and ketones?
 - A Acidified potassium dichromate solution
 - B 2,4-dinitrophenylhydrazine solution
 - C Fehling's solution
 - D Tollen's reagent



9 When ammonium chloride is dissolved in water the temperature decreases. Which one of the following shows the changes for ΔH and ΔS ?

 ΔH ΔS

- A negative negative
- B negative positive
- C positive negative
- D positive positive
- 10 cm³ of 6 M hydrochloric acid were diluted to form 0.5 M hydrochloric acid. Which one of the following is the volume of water to be added?
 - A 50 cm³

Rewarding L

Rewarding L

- B 110 cm³
- C 120 cm³
- D 290 cm³

[Turn over



Section B

Answer all six questions in this section

11 Write equations for the reaction of the following oxides and chlorides of the third period with water.

oxide/chloride	equation
sodium oxide	
sulfur trioxide	
chlorine(VII) oxide	
phosphorus(V) oxide	
phosphorus(V) chloride	

[5]

ng Learning

Rowardin

Rewardin

)

Rewardin

DED xg Learning

O.

DED va Lasmina

y Learning
Rewardin


Rewarding

The Control of the Contro

DE SI Lagraige

G:







		_ [2
(ii)	Explain how these hydrocarbon fuels increase the concentration of carbonic dioxide in the atmosphere.	on
		_ ['
(i)	Photosynthesis	
		[
(ii)	Respiration	
		[
	Stat diox (i)	State and explain the effect of the following on the concentration of carbon dioxide in the atmosphere.

ag Learning
Rewardin

Rowardin

Powerdin

Rowardin

Rowardin

Powerdin

Rowardin

Rowardin

Rowardin

Rowardin

Rowardin

Rowardin

Rewardin

Remarding
Sp Learning

Rewardin

Rowards,

Jaming

Rewardin

newardin Department

Rewardin

Rewardin 200 201 Learning



(c)	The concentration of carbon dioxide causes the "greenhouse effect". Explain what is meant by the "greenhouse effect".

[Turn over

10135

Œ

Rewarding L

Committee Com

Rewarding L

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

A Learning

GE Rewarding L

Rowarding L

Rewarding L

Rowarding L

Rowarding L


Rowarding L

Rowarding L

Rowarding L



- 13 Lithium hydride is produced by passing hydrogen gas over lithium. The reaction is complete at a temperature of 725 °C. The enthalpy of formation can be determined by using a Born–Haber cycle or Hess's law.
 - (a) An outline Born–Haber cycle with enthalpy values is shown below. However the enthalpy values have not been given + or signs.



(i) Write + or – before each enthalpy value on the diagram.

(ii) Calculate the molar enthalpy of formation for lithium hydride.

[2]
 4

[2]



	(iii)	The lattice e kJ mol ⁻¹ :	nthalpies fo	or the other Gr	oup I hydrides are	shown below in
		NaH	+864;	KH +729;	RbH +704;	CsH +656
		In terms of the	ne Born–Ha	aber cycle, sug	gest why these va	alues decrease.
						[2]
(b)				ation for lithium um hydride wit	3	etermined using the
			2Li + 2H ₂	O → 2LiOH	$+ H_2 \Delta H = -440$) kJ
			LiH + H ₂ C	→ LiOH +	$H_2 \Delta H = -130$) kJ
		culate the mo	lar enthalp	y of formation	of lithium hydride	using the enthalpy
						[2]
(c)				l – T∆S explai has a negative		ergy change for the
						[2]

Œ

Rewarding L

Rewarding L

Committee Com

Rewarding L

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

A Learning

GE Rewarding L

Rowarding L

Rewarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L

10135

[Turn over



(1)	Explain why the hydride ion is a nucleophile.
/::\	Draw the machanism for the reduction of propagate with hydride icae
(III)	Draw the mechanism for the reduction of propanone with hydride ions. The intermediate is reacted with water to form the organic product.
(iii)	Name the organic product from the reduction of propanone.
(iv)	Carboxylic acids are also reduced by lithal. Write an equation for the reduction of propanoic acid with lithal using [H] as the reducing agent.

ng Learning
Rewardin

Rowardin

Powerdin

Rowardin

Rowardin

Powerdin

Rowardin

Rowardin

Rowardin

Rowardin

Rowardin

Rowardin

Rewardin

Remarding
Sp Learning

Rewardin

Rewardin

Page 1 Learning

Page 2 Learning

Page 2 Learning

Page 3 Learning

Page 3 Learning

Page 3 Learning

Page 4 Learning

Page 5 Learning

Rewarding 19 Learning

Rewardin

newardir.

Rewardin

Rewardin 200 201 Learning



14	"su	anoethanoic acid, CH ₂ CNCOOH, is an important intermediate in the production perglue". It is a white solid with a melting point of 65–67 °C. It is a stronger acid n ethanoic acid.	
	(a)	Cyanoethanoic acid is prepared by the reaction of potassium chloroethanoate with potassium cyanide. The cyanoethanoic acid is liberated by the addition of hydrochloric acid.	
		Write equations for the two reactions taking place.	
			[2]
	(b)	Cyanoethanoic acid is very soluble in water and may be purified by recrystallisation. However, it cannot be distilled at atmospheric pressure since 160 °C it is decarboxylated to form ethanenitrile, CH ₃ CN.	e at
		(i) Explain how you would choose a suitable solvent to recrystallise cyanoethanoic acid.	
			[4]
		(ii) Write the equation for the decarboxylation of cyanoethanoic acid.	
			[1]
			ove

Œ

Rewarding L

Learning

Rewarding L.

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

A Learning

GE Rewarding L

Rowarding L



	(c)	has	anoethanoic acid can be esterified by reaction with alcohols. The methyl esternoethanoic acid can be esterified by reaction with alcohols. The methyl esternoethan a boiling point of 205–207°C and the ethyl esternas a boiling point of 210°C.	er
		(i)	Comment on these boiling points with regard to intermolecular forces.	
				_
				_
			[4]
		(ii)	Write an equation for the formation of the methyl ester	1]
		(iii)	Name a catalyst for the reaction.	1]
				.,
10135				

ag Learning
Rewardin

Rowardin

Powerdin

Rowardin

Rowardin

Powerdin

Rowardin

Rowardin

Rowardin

Rowardin

Rowardin

Rowardin

Rewardin

Remarding
Sp Learning

Rewardin

Revearded

To J. Learning

Rewardin

newardin Department

Rewardin

Rewardin 200 201 Learning



(d) Ethyl cyanoethanoate reacts with formaldehyde to give a cyanoacrylic ester which is the basis of "superglue".

$$\mathsf{CH}_2\!\!=\!\!\mathsf{O} \; + \; \mathsf{CH}_2\mathsf{CNCO}_2\mathsf{C}_2\mathsf{H}_5 \; \longrightarrow \; \mathsf{CH}_2\!\!=\!\!\mathsf{C}(\mathsf{CN})\mathsf{CO}_2\mathsf{C}_2\mathsf{H}_5 \; + \; \mathsf{H}_2\mathsf{O}$$

(i) Give the systematic name for formaldehyde.

_____[1]

(ii) Draw the structure of the cyanoacrylic ester, showing **all** the bonds present and explain whether it has E/Z isomers.

[3]

(iii) Draw the polymeric structure formed from the cyanoacrylic ester showing **three** repeating units.

[2]


[Turn over

10135

Rewarding L



(e) The titration curve for the reaction of cyanoethanoic acid with sodium hydroxide is shown below.



The cyanoethanoic acid reacts to form a salt. The pH of the solution is given by the equation:

$$pH = pK_a + log \frac{[salt]}{[acid]}$$



(i)	Write the equation for the dissociation constant, K_a , of cyanoethanoic acid.
(ii)	The concentration of the salt is the same as the concentration of the acid when the titration is half-complete. Use the titration curve to determine the pK $_{\rm a}$ and consequently the K $_{\rm a}$.
(iii	[2] Explain which indicator would be suitable for this titration.
	[2]
10135	[Turn over

Rewarding L

Rewarding L

Rewarding L

Company of the state of the sta

Learning L

Learning

Rowarding L

Rewarding L

I Learning

C. C. Rewarding I.

C. Rewarding I.

C. C. Rewarding I.

C. C. Rewarding I.

GE Rewarding L

Rowarding L

Rewarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L



15			fats react with potassium hydroxide and are saponified. The carboxylic om the oil/fat form potassium salts.	
	` ,	rour hydi con	saponification value of a fat was determined by adding 9.0 g of the fat to nd-bottomed flask containing 50.0 cm ³ of 1.00 M ethanolic potassium roxide and a few pieces of porous pot. The flask was fitted with a reflux denser and the mixture boiled for one hour by which time the solution war. The remaining solution was titrated with 0.50 M hydrochloric acid.	
		(i)	Suggest why ethanolic potassium hydroxide was used rather than aqueo potassium hydroxide.	ous
				[1]
		(ii)	Explain why the reaction took one hour and was not instantaneous.	
				[1]
		(iii)	Suggest the purpose of the porous pot.	
				[1]
10135				

Signaming

Rewarding

Signaming

Rewardin

Rewardin DED og Learning Rewardin DED ng Learning Rewardin ng Learning Rewardin ng Learnina Rewardin ng Learning G G DED ng Learning Rewardin newarding ag Learnino Rewardin DED ng Lesmina Rewardin DED ng Learning Rewardin ng Learning Rewardin DED og Learning Rewarding Ng Learning Rewardin

ewarding og Leaming Rewarding

newarding ng Learning Rewarding

Rewardin 200 201 Learning



		[1]
(v)	Calculate the saponification value of the fat if 15.0 cm ³ of the 0.50 M hydrochloric acid were required.	
		[3]

[Turn over

10135

Œ

Rowarding L

A Learning

Rewarding L

Rowarding L.

A Learning

GE Rewarding L

Rowarding L

Rewarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L



	(b)		e mole of an oil/fat, when hydrolysed, produced one mole of glycerol, one le of oleic acid, $\rm C_{18}H_{34}O_2$, and two moles of palmitic acid, $\rm C_{16}H_{32}O_2$.	
		(i)	Draw the possible structures of the oil/fat.	
				[3]
		(ii)	Explain whether any of the structures drawn have asymmetric centres.	
				[2]
10135				

Sy Learning
Rewarding
Sy Learning

Rewardin

Rewardin DED og Learning Rewardin DED og Learning Rewardin ng Learning Rewardin ng Learnina Rewardin ng Learning G G DED ng Learning Rewardin newarding ag Learnino Rewardin DED ng Lesmina Rewardin DED ng Learning Rewardin ng Learning Rewardin DED og Learning Rewarding Ng Learning Rewardin

Rewarding

ig Learning

Rewardin

DED og Learning
Rewardin

ng Learning
Rowarding
ng Learning

G.



(c)	igni	relative molecular mass, RMM, of a fatty acid can be determined by the tion of the silver salt of the fatty acid. A fatty acid suspected of being oleic was reacted with silver carbonate. The silver salt was heated to leave or er.	
	(i)	Write the equation for the reaction of the acid, RCOOH, with silver carbonate.	
			_ [1]
	(ii)	0.130 g of the pure silver salt, when heated to complete decomposition, gave 0.036 g of pure silver. Calculate the RMM of the silver salt.	
			[3]
	(iii)	Use the RMM calculated in part (ii) to confirm the identity of the fatty acid	.t

Œ

Rewarding L

A Learning

Rewarding L.

Rowarding L.
Rowarding L.
Rowarding L.
Rowarding L.
Rowarding L.
Rowarding L.
Rowarding L.
Rowarding L.
Rowarding L.

A Learning

GE Rewarding L

Rowarding L

Rewarding L

Rowarding L

10135

[Turn over



(c	d) Ara 5Z ste	achidonic acid, $C_{20}H_{32}O_2$, has four double bonds which are listed as $-8Z-11Z-14Z$. Suggest a structure of the acid showing the reochemistry about the double bonds.	
			[2]
(€	e) Ole	eic acid is insoluble in water but sodium oleate is soluble.	
	(i)	Explain why oleic acid is insoluble.	
			[1]
	(ii)	Explain why sodium oleate is soluble.	
			[2]
	(iii)	Explain whether a solution of sodium oleate is alkaline, acidic or neutral.	
			[2]
10135			

ng Learning
Rewardin

Rowardin

Powerdin

Rowardin

Rowardin

Powerdin

Rowardin

Rowardin

Rowardin

Rowardin

Rowardin

Rowardin

Rewardin

Remarding
Sp Learning

Rewardin

Revearded

To J. Learning

Rewardin

newardin Department

Rewardin

Rewardin 200 201 Learning



1	16	lodi	ine dissolves in cyclohexane but reacts reversibly with cyclohexene.
		(a)	Describe, giving experimental details, how you would use 100 cm ³ of cyclohexane to extract as much solid iodine as possible from a solution of iodine in water.
			[5]
			[5]
			Quality of written communication [2]
			[Turn over
1013	35		-

Rewarding L

Rewarding L

A Learning
Rewarding L

Rewarding L

Rewarding L

Rewarding L

Learning

L Learning

L Learning

L Learning

Rewarding L

Rewarding L

Rewarding L

Rowarding L

Rewarding L

I Learning

C. C. Rewarding I.

C. Rewarding I.

C. C. Rewarding I.

C. C. Rewarding I.

Rewarding L

Rowarding L

Rewarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L



(b) The equilibrium equation for the reaction of cyclohexene with iodine is:

ng Learning

E

E

G:

The equilibrium constant for the reaction at 25 $^{\circ}$ C is 20 mol dm⁻³ and at 35 $^{\circ}$ C is 13 mol dm⁻³.

(i)	Explain	whether	the	forward	reaction	is	endothermic of	or	exothermic
-----	---------	---------	-----	---------	----------	----	----------------	----	------------

			[2]
			1/1

(ii)	Suggest how you could determine the change in the concentration of I ₂	with
	time.	

		[3]
		[ა



	when equilibrium is reached?[1
(iv)	Calculate the equilibrium concentration of iodine, $\rm I_2$, in g dm $^{-3}$ in a system at equilibrium at 35 °C when the equilibrium concentration of $\rm C_6H_{10}I_2$ is 0.001 M and that of $\rm C_6H_{10}$ is 0.01 M.
c) Bro	[3] mine water is used to test for the presence of unsaturation. Suggest why
	ine is not used.
	THIS IS THE END OF THE QUESTION PAPER

Rewarding L

Rewarding L

Rewarding L

Company of the state of the sta

Rewarding L

Rewarding L

Learning

Rowarding L

GE Rewarding L

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

Rowarding Loaming

A Learning

GE Rewarding L

Rowarding L

Rewarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L

Rowarding L





ng Learning G. ng Learning G: ng Learning Rewardin ng Learnin-Rewards DED ig Learning Œ. DED va Leaming Rewardin **2** 0 GC:) G:

3


Page Searching
To Learning

ng Learning
Rowarding
Rowarding
Rowarding
Rowarding

ng Learning
Rewarding

Œ.





Rewarding I.

A Learning

Rowarding L



DO NOT WRITE ON THIS PAGE

For Examiner's use only				
Question Number	Marks			
Section A				
1–10				
Section B				
11				
12				
13				
14				
15				
16				

Ag Learning
Rowarding
Ag Learning
Rowarding
Rowarding

G:

E

)

Rowarding

Page Learning

Rowarding

E

y Learning

Rewardin

DE ng Learning

Rowarding

Rowarding

Rowarding

Rowarding

Rowarding

)

G.

ng Learning

Rewards

20 xg Learning

Rewarding ag Learning

Rewardin

G:

Total Marks

Examiner Number

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA will be happy to rectify any omissions of acknowledgement in future if notified.

