

Surname	Centre Number	Candidate Number
Other Names		0



**GCSE – NEW**

3430U50-1



S18-3430U50-1

**SCIENCE (Double Award)**

**Unit 5 – CHEMISTRY 2  
FOUNDATION TIER**

THURSDAY, 17 MAY 2018 – MORNING

1 hour 15 minutes

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1.	5	
2.	8	
3.	5	
4.	13	
5.	8	
6.	6	
7.	11	
8.	4	
<b>Total</b>	<b>60</b>	

3430U501  
01

**ADDITIONAL MATERIALS**

In addition to this examination paper you will need a calculator and a ruler.

**INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen. Do not use gel pen. Do not use correction fluid.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer **all** questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the additional page at the back of the booklet, taking care to number the question(s) correctly.

**INFORMATION FOR CANDIDATES**

The number of marks is given in brackets at the end of each question or part-question.

Question **6** is a quality of extended response (QER) question where your writing skills will be assessed.

The Periodic Table is printed on the back cover of this paper and the formulae for some common ions on the inside of the back cover.



MAY183430U50101

Answer all questions.

1. (a) Salts can be produced by reacting acids with alkalis.

(i) Complete the following equation for the reaction between an acid and alkali. [1]



(ii) **Circle** the word which best describes the reaction between an acid and an alkali. [1]

**displacement**


**neutralisation**

**oxidation**

**reduction**

(b) Indigestion can be caused by excess hydrochloric acid in the stomach. To treat indigestion, antacid powders are commonly used.

A group of pupils used the following apparatus to compare three brands of antacid powder, to see which was the most effective at treating acid indigestion.



They added an equal mass of each of the antacid powders to separate beakers, containing equal amounts of hydrochloric acid and universal indicator.

They stirred the mixture and recorded the time taken for the universal indicator to turn green in each beaker. They carried out the test three times for each antacid powder. Their results are shown in the table.



Antacid powder	Time taken for the universal indicator to turn green (min : s)			
	Result 1	Result 2	Result 3	Mean
Brand 1	5 : 25	5 : 36	5 : 14	5 : 25
Brand 2	4 : 28	3 : 20	4 : 32	4 : 30
Brand 3	2 : 28	2 : 30	2 : 44	2 : 34

- (i) State which **two** results were used to calculate the mean value for brand 2. [1]

..... and .....

- (ii) Convert the mean time for brand 2 into **seconds**. [1]

Mean time = ..... s

- (iii) Give the reason why the results suggest that brand 3 is the best powder for treating acid indigestion. [1]

.....

.....



2. (a) Chromium is one of the metals found in stainless steel. The equation shows how chromium is produced industrially by reacting chromium oxide with aluminium.



- (i) The reaction is highly exothermic.

Give the meaning of the term *exothermic*. [1]

- .....
- (ii) During the reaction, oxidation and reduction happens.

I. Name the substance which is oxidised. [1]

.....

II. State what is meant by *reduction*. [1]

- .....
- (iii) State what the equation tells you about the relative reactivities of chromium and aluminium. [1]

.....


.....



- (b) Copper is able to displace silver from a solution of silver nitrate. The equation for this reaction is given below.



A teacher demonstrated this reaction to her class. The photographs show the beaker before and after the reaction had taken place.



- (i) Explain how the changes show that this chemical reaction has taken place. [2]

.....


.....

.....

- (ii) Complete the symbol equation for the reaction by

- giving the formula for silver nitrate
- balancing the overall equation

[2]






8



3. (a) Magnesium reacts with chlorine to form magnesium chloride.

The following diagram shows the electronic structures of magnesium and chlorine atoms.




- (i) **Draw arrows on the diagram** to show how electrons are transferred between the magnesium and chlorine atoms during the formation of magnesium chloride. [1]
- (ii) **Complete the table** giving information about the charge and electronic structure of the magnesium and chloride **ions** that are formed. [2]

Ion	Charge	Electronic structure
magnesium	+2	.....
chloride	.....	2,8,8



(b) Chlorine gas,  $\text{Cl}_2$ , consists of two chlorine atoms bonded together.

(i) Draw a diagram in the box to show how the atoms bond to form a chlorine molecule. [1]



(ii) Give the name of this type of bonding.

.....

[1]



4. (a) Crude oil is a fossil fuel and is described as a non-renewable resource.

(i) Describe how crude oil was formed.

[2]

.....


.....

(ii) Give the meaning of the term *non-renewable*.

[1]

.....

(b) To make crude oil more useful, it is separated into fractions.



(i) Complete the following sentences.

[2]

Crude oil is separated into different fractions by a process called fractional

.....

The fractions can be separated because they have different

.....

(ii) **Circle** the word which best describes crude oil.

[1]

**mixture**

**element**

**compound**





(c) One of the fractions obtained from crude oil contains hexane,  $C_6H_{14}$ .

Calculate the percentage by mass of carbon in hexane.






[2]

$$A_r(H) = 1 \quad A_r(C) = 12$$

Percentage = ..... %



(d) The diagram shows the structures of five different hydrocarbons, **A-E**.

**A****B****C****D****E**

Use letters **A-E** in your answers to parts (i) and (ii).

(i) Give the structure that represents propene. .... [1]

(ii) Identify the structures that fit the following descriptions. [2]

Hydrocarbons with the general formula  $\text{C}_n\text{H}_{2n+2}$  .....

Unsaturated molecules .....



(e) Plastics are made from chemicals that are obtained from crude oil. Supermarkets in Wales were the first in the UK to charge their customers for plastic bags. This was to reduce the amount of plastic waste generated.

Give **two** methods of plastic waste disposal that lead to environmental problems. Explain the problem linked to each method. [2]

Method 1 .....

Problem .....

.....

Method 2 .....

Problem .....

.....

13



5. There are a number of factors that should be taken into consideration when deciding what makes the 'best fuel'.

Information was collected about various factors for three fuels, **A**, **B** and **C**.

**Fuel A**

- Existing supplies will last around 50 years
- Releases 2.8 kJ of energy per gram of fuel burned
- Costs 0.03p per gram of fuel burned
- Burns very easily and no storage issues
- Releases carbon dioxide and water vapour when it burns

**Fuel B**

- There are infinite supplies of this fuel
- Releases 44.1 kJ of energy per gram of fuel burned
- Costs 0.18p per gram of fuel burned
- Burns very easily but can be difficult to store
- Releases water vapour when it burns

**Fuel C**

- Existing supplies will last around 250 years
- Releases 1.2 kJ of energy per gram of fuel burned
- Costs 0.04p per gram of fuel burned
- Burns very easily and fairly easy to store
- Releases carbon dioxide, sulfur dioxide and water vapour when it burns

This information was analysed by a group of students to decide what they considered to be the 'best fuel'.



- (a) Give the reason why the information about how easily each fuel burns was not useful to the students when reaching their decision. [1]

.....

.....

- (b) One of the students based his decision purely on a judgement of the available supply of each fuel. Choose the order that shows his conclusion. Place a tick (✓) in the appropriate box. [1]

best fuel	<b>A</b>	<b>A</b>	<b>B</b>	<b>B</b>	<b>C</b>	<b>C</b>
↑	<b>B</b>	<b>C</b>	<b>A</b>	<b>C</b>	<b>A</b>	<b>B</b>
	<b>C</b>	<b>B</b>	<b>C</b>	<b>A</b>	<b>B</b>	<b>A</b>
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

- (c) Which of the following statements best describes how the fuels affect the environment when they burn? Tick (✓) the correct answer and give the reason for your choice. [2]

- all of the fuels contribute to acid rain and global warming when they burn
- fuels **A** and **C** contribute to acid rain and global warming when they burn
- only fuel **C** contributes to acid rain and global warming when it burns
- none of the fuels contribute to acid rain and global warming when they burn

Reason .....

.....



- (d) The cost efficiency of fuel **A** can be calculated as follows:

$$\text{cost efficiency} = \frac{2.8}{0.03} = 93.3 \text{ kJ/p}$$

Use the information given for fuel **B** and this example to calculate the cost efficiency of fuel **B**. [2]

Cost efficiency = ..... kJ/p

- (e) The students eventually agreed on the following rank order for the fuels.

best fuel




In the table below, tick (✓) **all** the statements that are **correct** and could therefore have been used in deciding upon this order. [2]


	(✓)
fuel <b>C</b> will run out after fuels <b>A</b> and <b>B</b>	<input type="checkbox"/>
fuel <b>C</b> is easier to store than fuel <b>A</b>	<input type="checkbox"/>
fuel <b>A</b> burns more easily than fuel <b>C</b>	<input type="checkbox"/>
fuel <b>B</b> is the cleanest fuel	<input type="checkbox"/>
fuel <b>B</b> is easier to store than fuel <b>C</b>	<input type="checkbox"/>
fuel <b>B</b> will never run out	<input type="checkbox"/>
fuel <b>A</b> is less harmful to the environment than fuel <b>C</b>	<input type="checkbox"/>
fuel <b>A</b> is less cost efficient than fuel <b>B</b>	<input type="checkbox"/>



6. The diagram shows the three stages used in the preparation of copper(II) sulfate crystals from copper(II) carbonate and sulfuric acid.



Stage 1



Stage 2



Stage 3

Describe and explain each stage of the preparation. Include an equation in your answer.

[6 QER]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



.....

6



7. A group of students carried out an investigation into the electrolysis of copper(II) sulfate solution. They used the apparatus shown to test the hypothesis:

***“the mass of copper that forms on the cathode increases as the time increases”***



To test the hypothesis, they weighed the cathode before placing it into the copper(II) sulfate solution and then again after allowing electrolysis to take place for varying times.


Their results are shown below.

Time (s)	Mass of copper formed (mg)		
	1	2	Mean
0	0	0	0
10	2.8	3.2	3.0
20	4.8	5.0	4.9
30	8.2	7.8	8.0
40	10.8	11.2	11.0
50	12.9	13.1	13.0
60	15.8	16.0	15.9





- (a) On the grid below, plot the mean mass of copper formed against time. Draw a suitable line. [3]



- (b) (i) Use the results collected at 30s and the following equation to calculate the percentage variation in these measurements. [2]

$$\text{percentage variation} = \frac{\text{furthest mass from the mean} - \text{mean mass}}{\text{mean mass}} \times 100$$


Percentage variation = ..... %

- (ii) The mass of copper formed is lower than expected. Give the most likely reason for this difference. [1]

.....  
.....  
.....



- (c) (i) Aluminium is extracted from molten aluminium oxide by electrolysis.




- I. Explain why aluminium forms at the cathode. [2]

.....

.....

.....

- II. Complete and balance the equation for the overall reaction that takes place. [2]




- (ii) Potassium can also be extracted through electrolysis of potassium carbonate.

Write the **formula** of potassium carbonate to complete the equation for the overall reaction. [1]



8. The burning of methane in air can be represented by the following equation.



The bond energies are given in the table below.

Bond	Bond energy (kJ)
C — H	413
O = O	498
O — H	464
C = O	805

- (a) Use the bond energy values to calculate the energy released when **all** the bonds in the carbon dioxide and water molecules are formed. [2]

Energy released = ..... kJ



- (b) The energy needed to break **all** the bonds in the methane and oxygen molecules is 2648 kJ.

Calculate the overall energy change for this reaction and use this value to explain why the reaction is exothermic. [2]

Overall energy change = ..... kJ

.....  
.....

4

**END OF PAPER**



Question number	<b>Additional page, if required. Write the question number(s) in the left-hand margin.</b>
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....

Examiner  
only



## FORMULAE FOR SOME COMMON IONS

POSITIVE IONS		NEGATIVE IONS	
Name	Formula	Name	Formula
aluminium	$\text{Al}^{3+}$	bromide	$\text{Br}^-$
ammonium	$\text{NH}_4^+$	carbonate	$\text{CO}_3^{2-}$
barium	$\text{Ba}^{2+}$	chloride	$\text{Cl}^-$
calcium	$\text{Ca}^{2+}$	fluoride	$\text{F}^-$
copper(II)	$\text{Cu}^{2+}$	hydroxide	$\text{OH}^-$
hydrogen	$\text{H}^+$	iodide	$\text{I}^-$
iron(II)	$\text{Fe}^{2+}$	nitrate	$\text{NO}_3^-$
iron(III)	$\text{Fe}^{3+}$	oxide	$\text{O}^{2-}$
lithium	$\text{Li}^+$	sulfate	$\text{SO}_4^{2-}$
magnesium	$\text{Mg}^{2+}$		
nickel	$\text{Ni}^{2+}$		
potassium	$\text{K}^+$		
silver	$\text{Ag}^+$		
sodium	$\text{Na}^+$		
zinc	$\text{Zn}^{2+}$		





24

# THE PERIODIC TABLE

2

1

Group

3

4

5

6

7

0

7 <b>Li</b> Lithium 3	9 <b>Be</b> Beryllium 4	11 <b>Na</b> Sodium 11	12 <b>C</b> Carbon 6	13 <b>Al</b> Aluminium 13	14 <b>N</b> Nitrogen 7	15 <b>O</b> Oxygen 8	16 <b>F</b> Fluorine 9	17 <b>Ne</b> Neon 10										
19 <b>K</b> Potassium 19	20 <b>Ca</b> Calcium 20	23 <b>Sc</b> Scandium 21	24 <b>Ti</b> Titanium 22	25 <b>V</b> Vanadium 23	26 <b>Cr</b> Chromium 24	27 <b>Mn</b> Manganese 25	28 <b>Fe</b> Iron 26	29 <b>Co</b> Cobalt 27	30 <b>Ni</b> Nickel 28	31 <b>Cu</b> Copper 29	32 <b>Zn</b> Zinc 30	35.5 <b>Ga</b> Gallium 31	36 <b>Ge</b> Germanium 32	37 <b>As</b> Arsenic 33	38 <b>Se</b> Selenium 34	39 <b>Br</b> Bromine 35	40 <b>Kr</b> Krypton 36	
37 <b>Rb</b> Rubidium 37	38 <b>Sr</b> Strontium 38	39 <b>Y</b> Yttrium 39	40 <b>Zr</b> Zirconium 40	41 <b>Nb</b> Niobium 41	42 <b>Mo</b> Molybdenum 42	43 <b>Tc</b> Technetium 43	44 <b>Ru</b> Ruthenium 44	45 <b>Rh</b> Rhodium 45	46 <b>Pd</b> Palladium 46	47 <b>Ag</b> Silver 47	48 <b>Cd</b> Cadmium 48	49 <b>In</b> Indium 49	50 <b>Sn</b> Tin 50	51 <b>Sb</b> Antimony 51	52 <b>Te</b> Tellurium 52	53 <b>I</b> Iodine 53	54 <b>Xe</b> Xenon 54	
55 <b>Cs</b> Caesium 55	56 <b>Ba</b> Barium 56	57 <b>La</b> Lanthanum 57	72 <b>Hf</b> Hafnium 72	73 <b>Ta</b> Tantalum 73	74 <b>W</b> Tungsten 74	75 <b>Re</b> Rhenium 75	76 <b>Os</b> Osmium 76	77 <b>Ir</b> Iridium 77	78 <b>Pt</b> Platinum 78	79 <b>Au</b> Gold 79	80 <b>Hg</b> Mercury 80	81 <b>Tl</b> Thallium 81	82 <b>Pb</b> Lead 82	83 <b>Bi</b> Bismuth 83	84 <b>Po</b> Polonium 84	85 <b>At</b> Astatine 85	86 <b>Rn</b> Radon 86	
87 <b>Fr</b> Francium 87	88 <b>Ra</b> Radium 88	89 <b>Ac</b> Actinium 89																

Key

relative atomic mass

$A_r$	Symbol Name Z
Symbol	
Z	

atomic number